

# A model for prediction of radon concentration in the Postojna cave

Asta Gregorič<sup>1,2</sup>

<sup>1</sup> Department of Environmental Sciences, Jožef Stefan Institute, Ljubljana, Slovenia

<sup>2</sup> Jožef Stefan International Postgraduate School (Ecotechnology, 2nd year)

asta.gregoric@ijs.si

Postojna Cave is the biggest of 12 show caves in Slovenia. Because of elevated radon concentrations, it has been under permanent radon survey since 1995 [1]. The driving force for air movement in the cave, and thus the inflow of fresh air and release of the cave air to the atmosphere, is the temperature difference between the cave air and outdoors [2]. In our study, we intend to predict radon concentration in the cave air on the basis of this temperature difference.

Measurements of radon concentration in the Postojna Cave were carried out continually (recorded once per hour) from July 2005 to October 2009 in the Great Mountain hall. Outdoor temperature has been recorded at the Postojna meteorological station and provided by the Environmental Agency of the Republic of Slovenia.

The cave behaves as a large chimney, and in the cold period the warmer radon-rich cave air is released to the colder outdoor atmosphere, allowing the inflow of fresh air with low radon levels. The radon levels in the cave are the highest when the outdoor temperature is similar to the cave temperature (10 °C) and, hence, the air movement is very low. Our calculations have shown that the effect of the difference between outdoor and cave temperature on radon concentration was delayed for three days, presumably because of the distance of the Great Mountain from the entrance (ca. 2 km). When daily outdoor temperature drops below – 6 °C, radon concentration grows higher resulting in an average value of  $644 \pm 235$  Bq m<sup>-3</sup>. It is assumed that ice and snow on the surface above the cave prevent the cave air from escaping the cave. In summer, the vertical air movement stops or is minimal and the horizontal air movement has the main influence on radon concentration in the cave.

For data analysis, the one dimensional transport equation provided by Nazaroff [3] has been used:

$\frac{dC}{dt} \approx -\frac{k_1|\Delta T|C}{L} - \lambda C + \Phi$  (1). Here,  $C$  is the radon concentration [Bq m<sup>-3</sup>],  $t$  is time [s],  $\Delta T$  is the difference

between outdoor temperature and cave temperature [K],  $L$  is the distance from the entrance to the Great Mountain [m],  $\lambda$  is radon decay constant ( $2.1 \times 10^{-6}$  s<sup>-1</sup>) and  $\Phi$  is radon source term [Bq m<sup>-3</sup> s<sup>-1</sup>]. Based on the temperature difference, this equation predicts radon concentration. In the analysis of the dataset from July 2005 to October 2007, the concentration reached its maximum (2,800 Bq m<sup>-3</sup>) in summer when the air draught in the cave was minimal or zero. Under these conditions, the radon source term of  $5.9 \times 10^{-3}$  Bq m<sup>-3</sup> s<sup>-1</sup> was obtained. The  $k$  values were calculated for the summer ( $k_1$ ) and winter regime ( $k_2$ ) separately, and the averages of  $0.19 \pm 1.07$  mm s<sup>-1</sup> K<sup>-1</sup> and  $0.93 \pm 31.13$  mm s<sup>-1</sup> K<sup>-1</sup>, respectively, were obtained.

The model, developed using the results of our previous long-term radon monitoring in the Postojna Cave, provides a relatively good prediction of radon concentration in the cave air, simply on the basis of the difference in air temperature in the cave and outdoors. Although successful, the model may certainly not replace the measurements but could reduce markedly their number, without diminishing the level of reliability of data needed for dose estimates for the personnel working in the cave.

References:

- [1] J. Vaupotič. Nanosize radon short-lived decay products in the air of the Postojna Cave. *Science of the Total Environment*, 393(1): 27-38, 2008.
- [2] J. Hakl, et al. Radon transport phenomena studied in Karst caves-international experiences on radon levels and exposures. *Radiation Measurements*, 28(1-6): 675-684, 1997.
- [3] W. W. Nazaroff. Radon transport from soil to air. *Reviews of Geophysics*, 30(2): 137-160, 1992.